

JOIN US FOR A LIVE EVENT



Learn From a Hematology and Oncology Expert

Get important insights on the efficacy and safety of REVLIMID for patients with newly diagnosed multiple myeloma

Live Q & A

Have your questions answered by an expert following the presentation



Speaker



Location



Date & Time

This is a promotional program, and no CME credits are offered.

Pursuant to the PhRMA Code on interactions with healthcare professionals, attendance at this promotional program is restricted to healthcare professionals. Accordingly, spouses and other guests who are not healthcare professionals may not attend this event.

Celgene will report transfers of value made to U.S. healthcare professionals pursuant to applicable law and company policy. To learn about how Celgene Corporation complies with the Physician Payments Sunshine Act, visit www.celgene.com/about/compliance/sunshine-act/.

The information you provide to Celgene will be used solely by Celgene and its agents in connection with this program. We do not sell or otherwise provide your personal information to third parties for marketing purposes.

Intended for specific recipient only; please do not forward or further disseminate.

You may also register by contacting Program Registration at pcaton@vectorhealth.net or calling (866) 991-3694. Please provide your name, degree, office address, phone number, and e-mail address. For questions, please contact your Celgene representative, at

Important Safety Information

WARNING: EMBRYO-FETAL TOXICITY, HEMATOLOGIC TOXICITY, and VENOUS and ARTERIAL THROMBOEMBOLISM

Embryo-Fetal Toxicity

Do not use REVLIMID during pregnancy. Lenalidomide, a thalidomide analogue, caused limb abnormalities in a developmental monkey study. Thalidomide is a known human teratogen that causes severe life-threatening human birth defects. If lenalidomide is used during pregnancy, it may cause birth defects or embryo-fetal death. In females of reproductive potential, obtain 2 negative pregnancy tests before starting REVLIMID treatment. Females of reproductive potential must use 2 forms of contraception or continuously abstain from heterosexual sex during and for 4 weeks after REVLIMID treatment. To avoid embryo-fetal exposure to lenalidomide, REVLIMID is only available through a restricted distribution program, the REVLIMID REMS® program.

Information about the REVLIMID REMS® program is available at www.celgeneriskmanagement.com or by calling the manufacturer's toll-free number 1-888-423-5436.

Hematologic Toxicity (Neutropenia and Thrombocytopenia)

REVLIMID can cause significant neutropenia and thrombocytopenia. Eighty percent of patients with del 5q MDS had to have a dose delay/reduction during the major study. Thirty-four percent of patients had to have a second dose delay/reduction. Grade 3 or 4 hematologic toxicity was seen in 80% of patients enrolled in the study. Patients on therapy for del 5q MDS should have their complete blood counts monitored weekly for the first 8 weeks of therapy and at least monthly thereafter. Patients may require dose interruption and/or reduction. Patients may require use of blood product support and/or growth factors.

Venous and Arterial Thromboembolism

REVLIMID has demonstrated a significantly increased risk of deep vein thrombosis (DVT) and pulmonary embolism (PE), as well as risk of myocardial infarction and stroke in patients with MM who were treated with REVLIMID and dexamethasone therapy. Monitor for and advise patients about signs and symptoms of thromboembolism. Advise patients to seek immediate medical care if they develop symptoms such as shortness of breath, chest pain, or arm or leg swelling. Thromboprophylaxis is recommended and the choice of regimen should be based on an assessment of the patient's underlying risks.

CONTRAINDICATIONS

Pregnancy: REVLIMID can cause fetal harm when administered to a pregnant female and is contraindicated in females who are pregnant. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential risk to the fetus

Severe Hypersensitivity Reactions: REVLIMID is contraindicated in patients who have demonstrated severe hypersensitivity (e.g., angioedema, Stevens-Johnson syndrome, toxic epidermal necrolysis) to lenalidomide

WARNINGS AND PRECAUTIONS

Embryo-Fetal Toxicity: See Boxed WARNINGS

- Females of Reproductive Potential: See Boxed WARNINGS**
- Males:** Lenalidomide is present in the semen of patients receiving the drug. Males must always use a latex or synthetic condom during any sexual contact with females of reproductive potential while taking REVLIMID and for up to 4 weeks after discontinuing REVLIMID, even if they have undergone a successful vasectomy. Male patients taking REVLIMID must not donate sperm
- Blood Donation:** Patients must not donate blood during treatment with REVLIMID and for 4 weeks following discontinuation of the drug because the blood might be given to a pregnant female patient whose fetus must not be exposed to REVLIMID

REVLIMID REMS® Program: See Boxed WARNINGS: Prescribers and pharmacies must be certified with the REVLIMID REMS program by enrolling and complying with the REMS requirements; pharmacies must only dispense to patients who are authorized to receive REVLIMID. Patients must sign a Patient-Physician Agreement Form and comply with REMS requirements; female patients of reproductive potential who are not pregnant must comply with the pregnancy testing and contraception requirements and males must comply with contraception requirements

Hematologic Toxicity: REVLIMID can cause significant neutropenia and thrombocytopenia. Monitor patients with neutropenia for signs of infection. Advise patients to observe for bleeding or bruising, especially with use of concomitant medications that may increase risk of bleeding. **MM:** Patients taking REVLIMID/dex or REVLIMID as maintenance therapy should have their complete blood counts (CBC) assessed every 7 days for the first 2 cycles, on days 1 and 15 of cycle 3, and every 28 days thereafter

Venous and Arterial Thromboembolism: See Boxed WARNINGS: Venous thromboembolic events (DVT and PE) and arterial thromboses (MI and CVA) are increased in patients treated with REVLIMID. Patients with known risk factors, including prior thrombosis, may be at greater risk and actions should be taken to try to minimize all modifiable factors (e.g., hyperlipidemia, hypertension, smoking). Thromboprophylaxis is recommended and the regimen should be based on patient's underlying risks. ESAs and estrogens may further increase the risk of thrombosis and their use should be based on a benefit-risk decision

Increased Mortality in Patients with CLL: In a clinical trial in the first-line treatment of patients with CLL, single agent REVLIMID therapy increased the risk of death as compared to single agent chlorambucil. Serious adverse cardiovascular reactions, including atrial fibrillation, myocardial infarction, and cardiac failure, occurred more frequently in the REVLIMID arm. REVLIMID is not indicated and not recommended for use in CLL outside of controlled clinical trials

Second Primary Malignancies (SPM): In clinical trials in patients with MM receiving REVLIMID, an increase of hematologic plus solid tumor SPM, notably AML and MDS, have been observed. Monitor patients for the development of SPM. Take into account both the potential benefit of REVLIMID and risk of SPM when considering treatment

Increased Mortality with Pembrolizumab: In clinical trials in patients with multiple myeloma, the addition of pembrolizumab to a thalidomide analogue plus dexamethasone resulted in increased mortality. Treatment of patients with multiple myeloma with a PD-1 or PD-L1 blocking antibody in combination with a thalidomide analogue plus dexamethasone is not recommended outside of controlled clinical trials

Hepatotoxicity: Hepatic failure, including fatal cases, has occurred in patients treated with REVLIMID/dex. Pre-existing viral liver disease, elevated baseline liver enzymes, and concomitant medications may be risk factors. Monitor liver enzymes periodically. Stop REVLIMID upon elevation of liver enzymes. After return to baseline values, treatment at a lower dose may be considered

Severe Cutaneous Reactions Including Hypersensitivity Reactions: Angioedema and severe cutaneous reactions including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported. DRESS may present with a cutaneous reaction (such as rash, or exfoliative dermatitis), eosinophilia, fever, and/or lymphadenopathy with systemic complications such as hepatitis, nephritis, pneumonitis, myocarditis, and/or pericarditis. These events can be fatal. Patients with a prior history of Grade 4 rash associated with thalidomide treatment should not receive REVLIMID. REVLIMID interruption or discontinuation should be considered for Grade 2-3 skin rash. REVLIMID must be discontinued for angioedema, Grade 4 rash, exfoliative or bullous rash, or if SJS, TEN, or DRESS is suspected and should not be resumed following discontinuation for these reactions

Tumor Lysis Syndrome (TLS): Fatal instances of TLS have been reported during treatment with lenalidomide. The patients at risk of TLS are those with high tumor burden prior to treatment. These patients should be monitored closely and appropriate precautions taken

Tumor Flare Reaction (TFR): TFR has occurred during investigational use of lenalidomide for CLL and lymphoma. Monitoring and evaluation for TFR is recommended in patients with MCL. Tumor flare may mimic the progression of disease (PD). In patients with Grade 3 or 4 TFR, it is recommended to withhold treatment with REVLIMID until TFR resolves to ≤Grade 1. REVLIMID may be continued in patients with Grade 1 and 2 TFR without interruption or modification, at the physician's discretion

Impaired Stem Cell Mobilization: A decrease in the number of CD34+ cells collected after treatment (>4 cycles) with REVLIMID has been reported. Consider early referral to transplant center to optimize timing of the stem cell collection

Thyroid Disorders: Both hypothyroidism and hyperthyroidism have been reported. Measure thyroid function before start of REVLIMID treatment and during therapy

Early Mortality in Patients with MCL: In another MCL study, there was an increase in early deaths (within 20 weeks), 12.9% in the REVLIMID arm versus 7.1% in the control arm. Risk factors for early deaths include high tumor burden, MIPI score at diagnosis, and high WBC at baseline (≥10 x 10⁹/L)

ADVERSE REACTIONS

Multiple Myeloma

- In newly diagnosed:** The most frequently reported Grade 3 or 4 reactions included neutropenia, anemia, thrombocytopenia, pneumonia, asthenia, fatigue, back pain, hypokalemia, rash, cataract, lymphopenia, dyspnea, DVT, hyperglycemia, and leukopenia. The highest frequency of infections occurred in Arm Rd Continuous (75%) compared to Arm MPT (56%). There were more Grade 3 and 4 and serious adverse reactions of infection in Arm Rd Continuous than either Arm MPT or Rd18
- The most common adverse reactions reported in ≥20% (Arm Rd Continuous): diarrhea (46%), anemia (44%), neutropenia (35%), fatigue (33%), back pain (32%), asthenia (28%), insomnia (28%), rash (26%), decreased appetite (23%), cough (23%), dyspnea (22%), pyrexia (21%), abdominal pain (21%), muscle spasms (20%), and thrombocytopenia (20%)
- Maintenance Therapy Post Auto-HSCT:** The most frequently reported Grade 3 or 4 reactions in ≥20% (REVLIMID arm) included neutropenia, thrombocytopenia, and leukopenia. The serious adverse reactions of lung infection and neutropenia (more than 4.5%) occurred in the REVLIMID arm
- The most frequently reported adverse reactions in ≥20% (REVLIMID arm) across both maintenance studies (Study 1, Study 2) were neutropenia (79%, 61%), thrombocytopenia (72%, 24%), leukopenia (23%, 32%), anemia (21%, 9%), upper respiratory tract infection (27%, 11%), bronchitis (5%, 47%), nasopharyngitis (2%, 35%), cough (10%, 27%), gastroenteritis (0%, 23%), diarrhea (55%, 39%), rash (32%, 8%), fatigue (23%, 11%), asthenia (0%, 30%), muscle spasm (0%, 33%), and pyrexia (8%, 21%)
- After at least one prior therapy:** The most common adverse reactions reported in ≥20% (REVLIMID/dex vs dex/placebo): fatigue (44% vs 42%), neutropenia (42% vs 6%), constipation (41% vs 21%), diarrhea (39% vs 27%), muscle cramp (33% vs 21%), anemia (31% vs 24%), pyrexia (28% vs 23%), peripheral edema (26% vs 21%), nausea (26% vs 21%), back pain (26% vs 19%), upper respiratory tract infection (25% vs 16%), dyspnea (24% vs 17%), dizziness (23% vs 17%), thrombocytopenia (22% vs 11%), rash (21% vs 9%), tremor (21% vs 7%), and weight decreased (20% vs 15%)

DRUG INTERACTIONS

Periodic monitoring of digoxin plasma levels is recommended due to increased C_{max} and AUC with concomitant REVLIMID therapy. Patients taking concomitant therapies such as erythropoietin stimulating agents or estrogen containing therapies may have an increased risk of thrombosis. It is not known whether there is an interaction between dex and warfarin. Close monitoring of PT and INR is recommended in patients with MM taking concomitant warfarin

USE IN SPECIFIC POPULATIONS

- PREGNANCY: See Boxed WARNINGS:** If pregnancy does occur during treatment, immediately discontinue the drug and refer patient to an obstetrician/gynecologist experienced in reproductive toxicity for further evaluation and counseling. There is a REVLIMID pregnancy exposure registry that monitors pregnancy outcomes in females exposed to REVLIMID during pregnancy as well as female partners of male patients who are exposed to REVLIMID. This registry is also used to understand the root cause for the pregnancy. Report any suspected fetal exposure to REVLIMID to the FDA via the MedWatch program at 1-800-FDA-1088 and also to Celgene Corporation at 1-888-423-5436
- LACTATION:** There is no information regarding the presence of lenalidomide in human milk, the effects of REVLIMID on the breastfed infant, or the effects of REVLIMID on milk production. Because many drugs are excreted in human milk and because of the potential for adverse reactions in breastfed infants from REVLIMID, advise female patients not to breastfeed during treatment with REVLIMID
- PEDIATRIC USE:** Safety and effectiveness have not been established in pediatric patients
- RENAL IMPAIRMENT:** Adjust the starting dose of REVLIMID based on creatinine clearance value and in patients on dialysis

Please see full [Prescribing Information](#), including Boxed WARNINGS.