

You are cordially invited to attend a live educational program:

DARZALEX[®]-BASED COMBINATION REGIMENS FOR MULTIPLE MYELOMA

PRESENTED BY

Rhonda Hewitt, MSN

Nurse Practitioner

Stanford Healthcare, Palo Alto, CA

**Thursday, June 13, 2019
6:00 PM**

LOCATION

Le Papillon

410 Saratoga Avenue, San Jose, CA 95129
(408) 296-3730

Please register with MedForce online at <http://www.medforcereg.net/SOMP/141232>
or with your Janssen Sales Representative, CATHERINE DUNCAN
at (610) 416-5451 by Thursday, June 06, 2019

In adherence with PhRMA guidelines, spouses or other guests are not permitted to attend company-sponsored programs.

For all attendees, please be advised that information such as your name and the value and purpose of any educational item, meal, or other items of value you receive may be publicly disclosed. If you are licensed in any state or other jurisdiction, or are an employee or contractor of any organization or governmental entity, that limits or prohibits meals from pharmaceutical companies, please identify yourself so that you (and we) are able to comply with such requirements.

Please note that the company prohibits the offering of gifts, gratuities, or meals to federal government employees/officials. Thank you for your cooperation.

This promotional educational activity is not accredited. The program content is developed by Janssen Biotech, Inc. Speakers present on behalf of the company and are required to present information in compliance with FDA requirements for communications about its medicines.

The personal information you provide may be used to contact you about your request to attend the Janssen Biotech, Inc., educational program using your preferred method of communication as indicated by you. This information may be shared with Janssen Biotech, Inc., its affiliates, and a third party for the sole purpose of completing your registration for this program and as required by law.

INDICATIONS

DARZALEX[®] (daratumumab) is indicated in combination with bortezomib, melphalan, and prednisone for the treatment of patients with newly diagnosed multiple myeloma who are ineligible for autologous stem cell transplant.

DARZALEX[®] is indicated in combination with lenalidomide and dexamethasone, or bortezomib and dexamethasone, for the treatment of patients with multiple myeloma who have received at least one prior therapy.

WARNINGS AND PRECAUTIONS

CONTRAINDICATIONS

DARZALEX[®] is contraindicated in patients with a history of severe hypersensitivity (eg, anaphylactic reactions) to daratumumab or any of the components of the formulation.

Please see Important Safety Information continued on the next page, including Warnings and Precautions for Infusion Reactions, Neutropenia, Thrombocytopenia, and Interference with Serological Testing and Determination of Complete Response.

Please CLICK HERE for the full Prescribing Information for DARZALEX[®].

IMPORTANT SAFETY INFORMATION (cont.)

WARNINGS AND PRECAUTIONS

Infusion Reactions – DARZALEX® can cause severe and/or serious infusion reactions, including anaphylactic reactions. In clinical trials, approximately half of all patients experienced an infusion reaction. Most infusion reactions occurred during the first infusion and were grade 1-2. Infusion reactions can also occur with subsequent infusions. Nearly all reactions occurred during infusion or within 4 hours of completing an infusion. Prior to the introduction of post-infusion medication in clinical trials, infusion reactions occurred up to 48 hours after infusion. Severe reactions have occurred, including bronchospasm, hypoxia, dyspnea, hypertension, laryngeal edema and pulmonary edema. Signs and symptoms may include respiratory symptoms, such as nasal congestion, cough, throat irritation, as well as chills, vomiting and nausea. Less common symptoms were wheezing, allergic rhinitis, pyrexia, chest discomfort, pruritus, and hypotension.

Pre-medicate patients with antihistamines, antipyretics, and corticosteroids. Frequently monitor patients during the entire infusion. Interrupt infusion for reactions of any severity and institute medical management as needed. Permanently discontinue therapy if an anaphylactic reaction or life-threatening (Grade 4) reaction occurs and institute appropriate emergency care. For patients with Grade 1, 2, or 3 reactions, reduce the infusion rate when re-starting the infusion.

To reduce the risk of delayed infusion reactions, administer oral corticosteroids to all patients following DARZALEX® infusions. Patients with a history of chronic obstructive pulmonary disease may require additional post-infusion medications to manage respiratory complications. Consider prescribing short- and long-acting bronchodilators and inhaled corticosteroids for patients with chronic obstructive pulmonary disease.

Interference with Serological Testing – Daratumumab binds to CD38 on red blood cells (RBCs) and results in a positive Indirect Antiglobulin Test (Indirect Coombs test). Daratumumab-mediated positive indirect antiglobulin test may persist for up to 6 months after the last daratumumab infusion.

Daratumumab bound to RBCs masks detection of antibodies to minor antigens in the patient's serum. The determination of a patient's ABO and Rh blood type are not impacted. Notify blood transfusion centers of this interference with serological testing and inform blood banks that a patient has received DARZALEX®. Type and screen patients prior to starting DARZALEX®.

Neutropenia – DARZALEX® may increase neutropenia induced by background therapy. Monitor complete blood cell counts periodically during treatment according to manufacturer's prescribing information for background therapies. Monitor patients with neutropenia for signs of infection. DARZALEX® dose delay may be required to allow recovery of neutrophils. No dose reduction of DARZALEX® is recommended. Consider supportive care with growth factors.

Thrombocytopenia – DARZALEX® may increase thrombocytopenia induced by background therapy. Monitor complete blood cell counts periodically during treatment according to manufacturer's prescribing information for background therapies. DARZALEX® dose delay may be required to allow recovery of platelets. No dose reduction of DARZALEX® is recommended. Consider supportive care with transfusions.

Interference with Determination of Complete Response – Daratumumab is a human IgG kappa monoclonal antibody that can be detected on both the serum protein electrophoresis (SPE) and immunofixation (IFE) assays used for the clinical monitoring of endogenous M-protein. This interference can impact the determination of complete response and of disease progression in some patients with IgG kappa myeloma protein.

Adverse Reactions – The most frequently reported adverse reactions (incidence $\geq 20\%$) in clinical trials were: infusion reactions, neutropenia, thrombocytopenia, fatigue, nausea, diarrhea, constipation, vomiting, muscle spasms, arthralgia, back pain, pyrexia, chills, dizziness, insomnia, cough, dyspnea, peripheral edema, peripheral sensory neuropathy and upper respiratory tract infection.

In patients who received DARZALEX® in combination with bortezomib, melphalan, and prednisone (DVMP), the most frequently reported adverse reactions (incidence $\geq 20\%$) were: upper respiratory tract infection (48%), infusion reactions (28%), and peripheral edema (21%). Serious adverse reactions ($\geq 2\%$ compared to the VMP arm) were pneumonia (11%), upper respiratory tract infection (5%), and pulmonary edema (2%). Treatment-emergent Grade 3-4 hematology laboratory abnormalities $\geq 20\%$ were lymphopenia (58%), neutropenia (44%), and thrombocytopenia (38%).

In patients who received DARZALEX® in combination with lenalidomide and dexamethasone, the most frequently reported adverse reactions (incidence $\geq 20\%$) were: upper respiratory tract infection (65%), infusion reactions (48%), diarrhea (43%), fatigue (35%), cough (30%), muscle spasms (26%), nausea (24%), dyspnea (21%) and pyrexia (20%). The overall incidence of serious adverse reactions was 49%. Serious adverse reactions ($\geq 2\%$ compared to Rd) were pneumonia (12%), upper respiratory tract infection (7%), influenza (3%), and pyrexia (3%). Treatment-emergent Grade 3-4 hematology laboratory abnormalities $\geq 20\%$ were neutropenia (53%) and lymphopenia (52%).

In patients who received DARZALEX® in combination with bortezomib and dexamethasone, the most frequently reported adverse reactions (incidence $\geq 20\%$) were: peripheral sensory neuropathy (47%), infusion reactions (45%), upper respiratory tract infection (44%), diarrhea (32%), cough (27%), peripheral edema (22%), and dyspnea (21%). The overall incidence of serious adverse reactions was 42%. Serious adverse reactions ($\geq 2\%$ compared to Vd) were upper respiratory tract infection (5%), diarrhea (2%) and atrial fibrillation (2%). Treatment-emergent Grade 3-4 hematology laboratory abnormalities $\geq 20\%$ were lymphopenia (48%) and thrombocytopenia (47%).

In patients who received DARZALEX® in combination with pomalidomide and dexamethasone, the most frequent adverse reactions ($>20\%$) were fatigue (50%), infusion reactions (50%), upper respiratory tract infection (50%), cough (43%), diarrhea (38%), constipation (33%), dyspnea (33%), nausea (30%), muscle spasms (26%), back pain (25%), pyrexia (25%), insomnia (23%), arthralgia (22%), dizziness (21%), and vomiting (21%). The overall incidence of serious adverse reactions was 49%. Serious adverse reactions reported in $\geq 5\%$ patients included pneumonia (7%). Treatment-emergent hematology Grade 3-4 laboratory abnormalities $\geq 20\%$ were anemia (30%), neutropenia (82%), and lymphopenia (71%).

In patients who received DARZALEX® as monotherapy, the most frequently reported adverse reactions (incidence $\geq 20\%$) were: infusion reactions (48%), fatigue (39%), nausea (27%), back pain (23%), pyrexia (21%), cough (21%), and upper respiratory tract infection (20%). The overall incidence of serious adverse reactions was 33%. The most frequent serious adverse reactions were pneumonia (6%), general physical health deterioration (3%), and pyrexia (3%). Treatment-emergent Grade 3-4 hematology laboratory abnormalities $\geq 20\%$ were lymphopenia (40%) and neutropenia (20%).

DRUG INTERACTIONS

Effect of Other Drugs on Daratumumab: The coadministration of lenalidomide, pomalidomide or bortezomib with DARZALEX® did not affect the pharmacokinetics of daratumumab.

Effect of Daratumumab on Other Drugs: The coadministration of DARZALEX® with bortezomib or pomalidomide did not affect the pharmacokinetics of bortezomib or pomalidomide.