

You are cordially invited to...

## KYPROLIS<sup>®</sup> (carfilzomib) +Darzalex<sup>®</sup> (daratumumab) and dexamethasone (DKd)

- Multiple myeloma disease state overview
- Mechanism of Action (MOA) of a second-generation Proteasome Inhibitor (PI) and CD38 Monoclonal Antibody (mAb)
- Review of the CANDOR study design
- Review of Progression-free Survival, Response Rates, and Minimal Residual Disease (MRD)
- Review of Safety Profile of the CANDOR study
- Communicate the important safety information for KYPROLIS<sup>®</sup>
- Review of the EQUULEUS Study

**Jennifer Knoche, RN, BSN, OCN, BMTCN**

Lead Practice Nurse

UCSF Hematology/BMT Ambulatory Care Center

**Wednesday, October 28, 2020**

**6:30 PM Pacific**

**Please RSVP To:**

**Divine Marcelo At (408) 838-0335 Or [dmarcelo@amgen.com](mailto:dmarcelo@amgen.com)**

### INDICATION

- KYPROLIS<sup>®</sup> (carfilzomib) is indicated in combination with dexamethasone or with lenalidomide plus dexamethasone or with daratumumab and dexamethasone for the treatment of adult patients with relapsed or refractory multiple myeloma who have received one to three lines of therapy.

### IMPORTANT SAFETY INFORMATION

#### Cardiac Toxicities

- New onset or worsening of pre-existing cardiac failure (e.g., congestive heart failure, pulmonary edema, decreased ejection fraction), cardiomyopathy, myocardial ischemia, and myocardial infarction including fatalities have occurred following administration of KYPROLIS. Some events occurred in patients with normal baseline ventricular function. Death due to cardiac arrest has occurred within one day of administration.
- Monitor patients for signs or symptoms of cardiac failure or ischemia. Evaluate promptly if cardiac toxicity is suspected. Withhold KYPROLIS for Grade 3 or 4 cardiac adverse reactions until recovery, and consider whether to restart at 1 dose level reduction based on a benefit/risk assessment.

**Please see additional Important Safety Information on the following pages.**



© 2020 Amgen Inc. All rights reserved.



# KYPROLIS<sup>®</sup> (carfilzomib) +Darzalex<sup>®</sup> (daratumumab) and dexamethasone (DKd)

## IMPORTANT SAFETY INFORMATION

### Cardiac Toxicities

- New onset or worsening of pre-existing cardiac failure (e.g., congestive heart failure, pulmonary edema, decreased ejection fraction), cardiomyopathy, myocardial ischemia, and myocardial infarction including fatalities have occurred following administration of KYPROLIS. Some events occurred in patients with normal baseline ventricular function. Death due to cardiac arrest has occurred within one day of administration.
- Monitor patients for signs or symptoms of cardiac failure or ischemia. Evaluate promptly if cardiac toxicity is suspected. Withhold KYPROLIS for Grade 3 or 4 cardiac adverse reactions until recovery, and consider whether to restart at 1 dose level reduction based on a benefit/risk assessment.
- While adequate hydration is required prior to each dose in Cycle 1, monitor all patients for evidence of volume overload, especially patients at risk for cardiac failure. Adjust total fluid intake as clinically appropriate.
- For patients  $\geq 75$  years, the risk of cardiac failure is increased. Patients with New York Heart Association Class III and IV heart failure, recent myocardial infarction, conduction abnormalities, angina, or arrhythmias may be at greater risk for cardiac complications and should have a comprehensive medical assessment prior to starting treatment with KYPROLIS and remain under close follow-up with fluid management.

### Acute Renal Failure

- Cases of acute renal failure, including some fatal renal failure events, and renal insufficiency (including renal failure) have occurred. Acute renal failure was reported more frequently in patients with advanced relapsed and refractory multiple myeloma who received KYPROLIS monotherapy. Monitor renal function with regular measurement of the serum creatinine and/or estimated creatinine clearance. Reduce or withhold dose as appropriate.

### Tumor Lysis Syndrome

- Cases of Tumor Lysis Syndrome (TLS), including fatal outcomes, have occurred. Patients with a high tumor burden should be considered at greater risk for TLS. Adequate hydration is required prior to each dose in Cycle 1, and in subsequent cycles as needed. Consider uric acid lowering drugs in patients at risk for TLS. Monitor for evidence of TLS during treatment and manage promptly, and withhold until resolved.

### Pulmonary Toxicity

- Acute Respiratory Distress Syndrome (ARDS), acute respiratory failure, and acute diffuse infiltrative pulmonary disease such as pneumonitis and interstitial lung disease have occurred. Some events have been fatal. In the event of drug-induced pulmonary toxicity, discontinue KYPROLIS.

### Pulmonary Hypertension

- Pulmonary arterial hypertension (PAH) was reported. Evaluate with cardiac imaging and/or other tests as indicated. Withhold KYPROLIS for PAH until resolved or returned to baseline and consider whether to restart based on a benefit/risk assessment.

### Dyspnea

- Dyspnea was reported in patients treated with KYPROLIS. Evaluate dyspnea to exclude cardiopulmonary conditions including cardiac failure and pulmonary syndromes. Stop KYPROLIS for Grade 3 or 4 dyspnea until resolved or returned to baseline. Consider whether to restart based on a benefit/risk assessment.

### Hypertension

- Hypertension, including hypertensive crisis and hypertensive emergency, has been observed, some fatal. Control hypertension prior to starting KYPROLIS. Monitor blood pressure regularly in all patients. If hypertension cannot be adequately controlled, withhold KYPROLIS and evaluate. Consider whether to restart based on a benefit/risk assessment.

Please see additional Important Safety Information on the following pages.

# KYPROLIS<sup>®</sup> (carfilzomib) +Darzalex<sup>®</sup> (daratumumab) and dexamethasone (DKd)

## IMPORTANT SAFETY INFORMATION (CONT'D)

### Venous Thrombosis

- Venous thromboembolic events (including deep venous thrombosis and pulmonary embolism) have been observed. Provide thromboprophylaxis for patients being treated with the combination of KYPROLIS with dexamethasone or with lenalidomide plus dexamethasone or with daratumumab and dexamethasone. The thromboprophylaxis regimen should be based on an assessment of the patient's underlying risks.
- For patients using hormonal contraception associated with a risk of thrombosis, consider an alternative method of effective contraception during treatment.

### Infusion-Related Reactions

- Infusion-related reactions, including life-threatening reactions, have occurred. Signs and symptoms include fever, chills, arthralgia, myalgia, facial flushing, facial edema, laryngeal edema, vomiting, weakness, shortness of breath, hypotension, syncope, chest tightness, or angina. These reactions can occur immediately following or up to 24 hours after administration. Premedicate with dexamethasone to reduce the incidence and severity of infusion-related reactions.

### Hemorrhage

- Fatal or serious cases of hemorrhage have been reported. Hemorrhagic events have included gastrointestinal, pulmonary, and intracranial hemorrhage and epistaxis. Promptly evaluate signs and symptoms of blood loss. Reduce or withhold dose as appropriate.

### Thrombocytopenia

- KYPROLIS causes thrombocytopenia with recovery to baseline platelet count usually by the start of the next cycle. Monitor platelet counts frequently during treatment. Reduce or withhold dose as appropriate.

### Hepatic Toxicity and Hepatic Failure

- Cases of hepatic failure, including fatal cases, have occurred. KYPROLIS can cause increased serum transaminases. Monitor liver enzymes regularly regardless of baseline values. Reduce or withhold dose as appropriate.

### Thrombotic Microangiopathy

- Cases of thrombotic microangiopathy, including thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS), including fatal outcome have occurred. Monitor for signs and symptoms of TTP/HUS. Discontinue if diagnosis is suspected. If the diagnosis of TTP/HUS is excluded, KYPROLIS may be restarted. The safety of reinitiating KYPROLIS is not known.

### Posterior Reversible Encephalopathy Syndrome (PRES)

- Cases of PRES have occurred in patients receiving KYPROLIS. If PRES is suspected, discontinue and evaluate with appropriate imaging. The safety of reinitiating KYPROLIS is not known.

### Progressive Multifocal Leukoencephalopathy (PML)

- Cases of PML, including fatal cases, have occurred. In addition to KYPROLIS, other contributory factors may include prior or concurrent use of immunosuppressive therapy. Consider PML in any patient with new onset of or changes in pre-existing neurological signs or symptoms. If PML is suspected, discontinue and initiate evaluation for PML including neurology consultation.

Please see additional Important Safety Information on the following pages.

# KYPROLIS<sup>®</sup> (carfilzomib) +Darzalex<sup>®</sup> (daratumumab) and dexamethasone (DKd)

## IMPORTANT SAFETY INFORMATION (CONT'D)

### Increased Fatal and Serious Toxicities in Combination with Melphalan and Prednisone in Newly Diagnosed Transplant-ineligible Patients

- In a clinical trial of transplant-ineligible patients with newly diagnosed multiple myeloma comparing KYPROLIS, melphalan, and prednisone (KMP) vs bortezomib, melphalan, and prednisone (VMP), a higher incidence of serious and fatal adverse reactions was observed in patients in the KMP arm. KMP is not indicated for transplant-ineligible patients with newly diagnosed multiple myeloma.

### Embryo-fetal Toxicity

- KYPROLIS can cause fetal harm when administered to a pregnant woman.
- Advise pregnant women of the potential risk to a fetus. Females of reproductive potential should use effective contraception during treatment with KYPROLIS and for 6 months following the final dose. Males of reproductive potential should use effective contraception during treatment with KYPROLIS and for 3 months following the final dose.

### Adverse Reactions

- The most common adverse reactions in the combination therapy trials: anemia, diarrhea, fatigue, hypertension, pyrexia, upper respiratory tract infection, thrombocytopenia, cough, dyspnea, and insomnia.

Notice: This event is conducted in accordance with the PhRMA Code on Interaction with Healthcare Professionals and is limited to invited healthcare professionals. Attendance by guests or spouse is not appropriate. Government employees are subject to state and federal laws and ethics rules that may limit your ability to receive any gifts, including meals, from pharmaceutical companies. If you are a state or federal employee, it is your responsibility to seek guidance and prior approval from your employer or site ethics counselor to attend this event.

State Laws: To comply with law and Amgen policies, Amgen is unable to offer food and beverages to (1) individuals with prescribing authority in Vermont or Minnesota; or (2) individuals employed by prescribers in Vermont who support the provision of healthcare. Please confirm the value of the meal with your Amgen representative before accepting the meal. You have the opportunity to opt-out of the meal and/or purchase your own meal, if applicable.

Please note that Amgen exercises diligence in reviewing the licensure of attendees and asks that you cooperate by disclosing all licensures in the sign-in/registration process. We appreciate your understanding and support.

Disclosure by Amgen: Amgen reports payments and transfers of value made to healthcare professionals and other healthcare related entities in accordance with federal and state laws, regulations and other transparency obligations. Any items of value provided by Amgen at this event (including the provision of meals and refreshments) may be subject to public disclosure. If you have questions regarding this matter please contact Amgen at 805-447-7422 or HCCSpendInquiry@amgen.com.

Please [click here](#) for the full Prescribing Information.

