

You are cordially invited to attend

Discover the Power of 1L RYBREVANT® + LAZCLUZE™ in EGFR+* Locally Advanced or mNSCLC

*Ex19del/L858R.

EGFR, epidermal growth factor receptor; ex19del, exon 19 deletions; L858R, exon 21 L858R substitution mutations; mNSCLC, metastatic non-small cell lung cancer.

PRESENTED BY

Robert Weber, MD
Director
St Mary's Medical Center Cancer Program

Speaker is a paid consultant of Johnson & Johnson

DATE

Wednesday, June 11, 2025
at 6:00 PM Pacific Time

LOCATION

Papillon Restaurant
37296 Mission Boulevard,
Fremont, California 94536
(510) 793-6331

We look forward to your participation in this informative discussion. Reserve your spot now by contacting your Johnson & Johnson Representative, Mark Hanes at mhanes2@its.jnj.com, (831) 277-3848.

EDUCATIONAL POINTS



Overview of mechanisms of action



Summary of indications, clinical trial design, and efficacy data



Safety profile and proactive therapy management

This is a non-CME educational program.

INDICATIONS

RYBREVANT® (amivantamab-vmjw) is indicated:

- in combination with LAZCLUZE™ (lazertinib) for the first-line treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations, as detected by an FDA-approved test.
- in combination with carboplatin and pemetrexed for the treatment of adult patients with locally advanced or metastatic NSCLC with EGFR exon 19 deletions or exon 21 L858R substitution mutations, whose disease has progressed on or after treatment with an EGFR tyrosine kinase inhibitor.

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Infusion-Related Reactions

RYBREVANT® can cause infusion-related reactions (IRR); signs and symptoms of IRR include dyspnea, flushing, fever, chills, nausea, chest discomfort, hypotension, and vomiting. The median time to IRR onset is approximately 1 hour.

Please see additional Important Safety Information continued on next page. Please read accompanying full Prescribing Information for RYBREVANT® and LAZCLUZE™.

In adherence with PhRMA guidelines, spouses or other guests are not permitted to attend company-sponsored programs.

For all attendees, please be advised that information such as your name and the value and purpose of any educational item, meal, or other items of value you receive may be publicly disclosed. If you are licensed in any state or other jurisdiction, or are an employee or contractor of any organization or governmental entity that limits or prohibits meals from pharmaceutical companies, please identify yourself so that you (and we) are able to comply with such requirements.

Please note that the company prohibits the offering of gifts, gratuities, or meals to federal government employees/officials. Thank you for your cooperation.

This promotional educational activity is not accredited. The program content is developed by Janssen Biotech, Inc. Speakers are paid consultants presenting on behalf of the company and are required to present information in compliance with FDA requirements for communications about its medicines.

The personal information you provide may be used to contact you about your request to attend the Janssen Biotech, Inc., educational program using your preferred method of communication as indicated by you. This information may be shared with Janssen Biotech, Inc., its affiliates, and a third party for the sole purpose of completing your registration for this program and as required by law.

IMPORTANT SAFETY INFORMATION (cont.)

RYBREVANT® with LAZCLUZE™

RYBREVANT® in combination with LAZCLUZE™ can cause infusion-related reactions. In MARIPOSA (n=421), IRRs occurred in 63% of patients treated with RYBREVANT® in combination with LAZCLUZE™, including Grade 3 in 5% and Grade 4 in 1% of patients. The incidence of infusion modifications due to IRR was 54% of patients, and IRRs leading to dose reduction of RYBREVANT® occurred in 0.7% of patients. Infusion-related reactions leading to permanent discontinuation of RYBREVANT® occurred in 4.5% of patients receiving RYBREVANT® in combination with LAZCLUZE™.

RYBREVANT® with Carboplatin and Pemetrexed

Based on the pooled safety population (n=281), IRR occurred in 50% of patients treated with RYBREVANT® in combination with carboplatin and pemetrexed, including Grade 3 (3.2%) adverse reactions. The incidence of infusion modifications due to IRR was 46%, and 2.8% of patients permanently discontinued RYBREVANT® due to IRR.

Premedicate with antihistamines, antipyretics, and glucocorticoids and infuse RYBREVANT® as recommended. Administer RYBREVANT® via a peripheral line on Week 1 and Week 2 to reduce the risk of infusion-related reactions. Monitor patients for signs and symptoms of infusion reactions during RYBREVANT® infusion in a setting where cardiopulmonary resuscitation medication and equipment are available. Interrupt infusion if IRR is suspected. Reduce the infusion rate or permanently discontinue RYBREVANT® based on severity.

Interstitial Lung Disease/Pneumonitis

RYBREVANT® can cause severe and fatal interstitial lung disease (ILD)/pneumonitis.

RYBREVANT® with LAZCLUZE™

In MARIPOSA, ILD/pneumonitis occurred in 3.1% of patients treated with RYBREVANT® in combination with LAZCLUZE™, including Grade 3 in 1.0% and Grade 4 in 0.2% of patients. There was one fatal

case (0.2%) of ILD/pneumonitis and 2.9% of patients permanently discontinued RYBREVANT® and LAZCLUZE™ due to ILD/pneumonitis.

RYBREVANT® with Carboplatin and Pemetrexed

Based on the pooled safety population, ILD/pneumonitis occurred in 2.1% treated with RYBREVANT® in combination with carboplatin and pemetrexed with 1.8% of patients experiencing Grade 3 ILD/pneumonitis. 2.1% discontinued RYBREVANT® due to ILD/pneumonitis.

Monitor patients for new or worsening symptoms indicative of ILD/pneumonitis (e.g., dyspnea, cough, fever). For patients receiving RYBREVANT® in combination with LAZCLUZE™, immediately withhold both drugs in patients with suspected ILD/pneumonitis and permanently discontinue if ILD/pneumonitis is confirmed. For patients receiving RYBREVANT® as a single agent or in combination with carboplatin and pemetrexed, immediately withhold RYBREVANT® in patients with suspected ILD/pneumonitis and permanently discontinue if ILD/pneumonitis is confirmed.

Venous Thromboembolic (VTE) Events with Concomitant Use of RYBREVANT® and LAZCLUZE™

RYBREVANT® in combination with LAZCLUZE™ can cause serious and fatal venous thromboembolic (VTE) events, including deep vein thrombosis and pulmonary embolism. The majority of these events occurred during the first four months of therapy.

In MARIPOSA, VTEs occurred in 36% of patients receiving RYBREVANT® in combination with LAZCLUZE™, including Grade 3 in 10% and Grade 4 in 0.5% of patients. On-study VTEs occurred in 1.2% of patients (n=5) while receiving anticoagulation therapy. There were two fatal cases of VTE (0.5%), 9% of patients had VTE leading to dose interruptions of RYBREVANT®, and 7% of patients had VTE leading to dose interruptions of LAZCLUZE™; 1% of patients had VTE leading to dose reductions of RYBREVANT®, and 0.5% of patients had VTE leading to dose reductions of LAZCLUZE™; 3.1% of patients had VTE leading to permanent discontinuation of RYBREVANT®, and 1.9% of patients had VTE leading to

Please see additional Important Safety Information continued on next page. Please read accompanying full Prescribing Information for RYBREVANT® and LAZCLUZE™.

IMPORTANT SAFETY INFORMATION (cont.)

permanent discontinuation of LAZCLUZE™. The median time to onset of VTEs was 84 days (range: 6 to 777).

Administer prophylactic anticoagulation for the first four months of treatment. The use of Vitamin K antagonists is not recommended. Monitor for signs and symptoms of VTE events and treat as medically appropriate.

Withhold RYBREVANT® and LAZCLUZE™ based on severity. Once anticoagulant treatment has been initiated, resume RYBREVANT® and LAZCLUZE™ at the same dose level at the discretion of the healthcare provider. In the event of VTE recurrence despite therapeutic anticoagulation, permanently discontinue RYBREVANT® and continue treatment with LAZCLUZE™ at the same dose level at the discretion of the healthcare provider.

Dermatologic Adverse Reactions

RYBREVANT® can cause severe rash including toxic epidermal necrolysis (TEN), dermatitis acneiform, pruritus, and dry skin.

RYBREVANT® with LAZCLUZE™

In MARIPOSA, rash occurred in 86% of patients treated with RYBREVANT® in combination with LAZCLUZE™, including Grade 3 in 26% of patients. The median time to onset of rash was 14 days (range: 1 to 556 days). Rash leading to dose interruptions occurred in 37% of patients for RYBREVANT® and 30% for LAZCLUZE™, rash leading to dose reductions occurred in 23% of patients for RYBREVANT® and 19% for LAZCLUZE™, and rash leading to permanent discontinuation occurred in 5% of patients for RYBREVANT® and 1.7% for LAZCLUZE™.

RYBREVANT® with Carboplatin and Pemetrexed

Based on the pooled safety population, rash occurred in 82% of patients treated with RYBREVANT® in combination with carboplatin and pemetrexed, including Grade 3 (15%) adverse reactions. Rash leading to dose reductions occurred in 14% of patients, and 2.5% permanently discontinued RYBREVANT® and 3.1% discontinued pemetrexed.

Instruct patients to limit sun exposure during and for 2 months after treatment with RYBREVANT® or LAZCLUZE™ in combination with RYBREVANT®. Advise patients to wear protective clothing and use broad-spectrum UVA/UVB sunscreen. Alcohol-free (e.g., isopropanol-free, ethanol-free) emollient cream is recommended for dry skin.

When initiating RYBREVANT® treatment with or without LAZCLUZE™, administer alcohol-free emollient cream to reduce the risk of dermatologic adverse reactions. Consider prophylactic measures (e.g. use of oral antibiotics) to reduce the risk of dermatologic reactions. If skin reactions develop, start topical corticosteroids

and topical and/or oral antibiotics. For Grade 3 reactions, add oral steroids and consider dermatologic consultation. Promptly refer patients presenting with severe rash, atypical appearance or distribution, or lack of improvement within 2 weeks to a dermatologist. For patients receiving RYBREVANT® in combination with LAZCLUZE™, withhold, reduce the dose, or permanently discontinue both drugs based on severity. For patients receiving RYBREVANT® as a single agent or in combination with carboplatin and pemetrexed, withhold, dose reduce or permanently discontinue RYBREVANT® based on severity.

Ocular Toxicity

RYBREVANT® can cause ocular toxicity including keratitis, blepharitis, dry eye symptoms, conjunctival redness, blurred vision, visual impairment, ocular itching, eye pruritus, and uveitis.

RYBREVANT® with LAZCLUZE™

In MARIPOSA, ocular toxicity occurred in 16% of patients treated with RYBREVANT® in combination with LAZCLUZE™, including Grade 3 or 4 ocular toxicity in 0.7% of patients. Withhold, reduce the dose, or permanently discontinue RYBREVANT® and continue LAZCLUZE™ based on severity.

RYBREVANT® with Carboplatin and Pemetrexed

Based on the pooled safety population, ocular toxicity occurred in 16% of patients treated with RYBREVANT® in combination with carboplatin and pemetrexed. All events were Grade 1 or 2.

Promptly refer patients with new or worsening eye symptoms to an ophthalmologist. Withhold, reduce the dose, or permanently discontinue RYBREVANT® based on severity.

Embryo-Fetal Toxicity

Based on its mechanism of action and findings from animal models, RYBREVANT® and LAZCLUZE™ can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential of the potential risk to the fetus.

Advise female patients of reproductive potential to use effective contraception during treatment and for 3 months after the last dose of RYBREVANT®.

Advise females of reproductive potential to use effective contraception during treatment with LAZCLUZE™ and for 3 weeks after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with LAZCLUZE™ and for 3 weeks after the last dose.

Adverse Reactions

RYBREVANT® with LAZCLUZE™

For the 421 patients in the MARIPOSA clinical trial who received RYBREVANT® in combination with LAZCLUZE™,

Please see additional Important Safety Information continued on next page. Please read accompanying full Prescribing Information for RYBREVANT® and LAZCLUZE™.

IMPORTANT SAFETY INFORMATION (cont.)

the most common adverse reactions ($\geq 20\%$) were rash (86%), nail toxicity (71%), infusion-related reactions (RYBREVANT[®], 63%), musculoskeletal pain (47%), stomatitis (43%), edema (43%), VTE (36%), paresthesia (35%), fatigue (32%), diarrhea (31%), constipation (29%), COVID-19 (26%), hemorrhage (25%), dry skin (25%), decreased appetite (24%), pruritus (24%), nausea (21%), and ocular toxicity (16%). The most common Grade 3 or 4 laboratory abnormalities ($\geq 2\%$) were decreased albumin (8%), decreased sodium (7%), increased ALT (7%), decreased potassium (5%), decreased hemoglobin (3.8%), increased AST (3.8%), increased GGT (2.6%), and increased magnesium (2.6%).

Serious adverse reactions occurred in 49% of patients who received RYBREVANT[®] in combination with LAZCLUZE[™]. Serious adverse reactions occurring in $\geq 2\%$ of patients included VTE (11%), pneumonia (4%), ILD/pneumonitis and rash (2.9% each), COVID-19 (2.4%), and pleural effusion and infusion-related reaction (RYBREVANT[®]) (2.1% each). Fatal adverse reactions occurred in 7% of patients who received RYBREVANT[®] in combination with LAZCLUZE[™] due to death not otherwise specified (1.2%); sepsis and respiratory failure (1% each); pneumonia, myocardial infarction, and sudden death (0.7% each); cerebral infarction, pulmonary embolism (PE), and COVID-19 infection (0.5% each); and ILD/pneumonitis, acute respiratory distress syndrome (ARDS), and cardiopulmonary arrest (0.2% each).

RYBREVANT[®] with Carboplatin and Pemetrexed

For the 130 patients in the MARIPOSA-2 clinical trial who received RYBREVANT[®] in combination with carboplatin and pemetrexed, the most common adverse reactions ($\geq 20\%$) were rash (72%), infusion-related reactions (59%), fatigue (51%), nail toxicity (45%), nausea (45%), constipation (39%), edema (36%), stomatitis (35%), decreased appetite (31%), musculoskeletal pain (30%), vomiting (25%), and COVID-19 (21%). The most common Grade 3 to 4 laboratory abnormalities ($\geq 2\%$) were decreased neutrophils (49%), decreased white blood cells (42%), decreased lymphocytes (28%), decreased platelets (17%), decreased hemoglobin (12%), decreased potassium (11%), decreased sodium (11%), increased alanine aminotransferase (3.9%), decreased albumin (3.8%), and increased gamma-glutamyl transferase (3.1%).

In MARIPOSA-2, serious adverse reactions occurred in 32% of patients who received RYBREVANT[®] in combination with carboplatin and pemetrexed. Serious adverse reactions in $>2\%$ of patients included dyspnea (3.1%), thrombocytopenia (3.1%), sepsis (2.3%), and pulmonary embolism (2.3%). Fatal adverse reactions occurred in 2.3% of patients who received RYBREVANT[®] in combination with carboplatin and pemetrexed; these included respiratory failure, sepsis, and ventricular fibrillation (0.8% each).

LAZCLUZE[™] Drug Interactions

Avoid concomitant use of LAZCLUZE[™] with strong and moderate CYP3A4 inducers. Consider an alternate concomitant medication with no potential to induce CYP3A4.

Monitor for adverse reactions associated with a CYP3A4 or BCRP substrate where minimal concentration changes may lead to serious adverse reactions, as recommended in the approved product labeling for the CYP3A4 or BCRP substrate.

cp-477344v1

Please read accompanying full Prescribing Information for RYBREVANT[®].

Please read accompanying full Prescribing Information for LAZCLUZE[™].